

# HIV Sequence Compendium 2017

## Editors

Brian Foley  
Los Alamos National Laboratory

Thomas Leitner  
Los Alamos National Laboratory

Cristian Apetrei  
University of Pittsburgh

Beatrice Hahn  
University of Pennsylvania

Ilene Mizrahi  
National Center for Biotechnology  
Information

James Mullins  
University of Washington

Andrew Rambaut  
University of Edinburgh

Steven Wolinsky  
Northwestern University

Bette Korber  
Los Alamos National Laboratory

## Contracting Officer's Representative

Anjali Singh  
Division of AIDS  
National Institute of Allergy and Infectious Diseases

## Los Alamos HIV Sequence Database and Analysis Staff

Werner Abfalterer, Elena E. Giori, Will Fischer, Peter Hraber,  
Jennifer Macke, James J. Szinger, Kshitij Wagh, Hyejin Yoon

This publication is funded by the Division of AIDS, National Institute of Allergy and Infectious Diseases, through interagency agreement AAI 12007-0000-01000 "HIV/SIV Database and Analysis Unit" with the U.S. Department of Energy.

Published by  
Theoretical Biology and Biophysics  
Group T-6, Mail Stop K710  
Los Alamos National Laboratory  
Los Alamos, New Mexico 87545 U.S.A.

LA-UR-17-25240

<https://www.hiv.lanl.gov/>



## HIV Sequence Compendium 2017: Preface

Published by  
Theoretical Biology and Biophysics  
Group T-6, Mail Stop K710  
Los Alamos National Laboratory  
Los Alamos, New Mexico 87545 U.S.A.

LA-UR-17-25240  
Approved for public release; distribution is unlimited.

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Los Alamos National Security, LLC, for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396.

This report was prepared as an account of work sponsored by an agency of the U.S. Government. Neither Los Alamos National Security, LLC, the U.S. Government nor any agency thereof, nor any of their employees make any warranty, express or implied, or assume any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represent that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by Los Alamos National Security, LLC, the U.S. Government, or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of Los Alamos National Security, LLC, the U.S. Government, or any agency thereof.

Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

This report was prepared as an account of work sponsored by NIH/NIAID/DAIDS under contract number AAI 12007-0000-01000 "HIV/SIV Database and Analysis Unit".

# Contents

<b>Contents</b>	<b>iii</b>
<b>I Preface</b>	<b>1</b>
I-1 Introduction . . . . .	1
I-2 Acknowledgements . . . . .	1
I-3 Citing the compendium or database . . . . .	1
I-4 About the PDF . . . . .	1
I-5 Genome maps . . . . .	2
I-6 HIV/SIV proteins . . . . .	3
I-7 Landmarks of the genome . . . . .	4
I-8 Amino acid codes . . . . .	6
I-9 Nucleic acid codes . . . . .	6
<b>II HIV-1/SIVcpz Complete Genomes</b>	<b>7</b>
II-1 Introduction . . . . .	7
II-2 Annotated features . . . . .	8
II-3 Sequences . . . . .	11
II-4 Alignments . . . . .	18
<b>III HIV-2/SIV Complete Genomes</b>	<b>155</b>
III-1 Introduction . . . . .	155
III-2 Annotated features . . . . .	156
III-3 Sequences . . . . .	158
III-4 Alignments . . . . .	161
<b>IV PLV Complete Genomes</b>	<b>227</b>
IV-1 Introduction . . . . .	227
IV-2 Sequences . . . . .	228
IV-3 Alignments . . . . .	231
<b>V HIV-1/SIVcpz Proteins</b>	<b>313</b>
V-1 Introduction . . . . .	313
V-2 Annotated features . . . . .	314
V-3 Sequences . . . . .	316
V-4 Alignments . . . . .	324
<b>VI HIV-2/SIV Proteins</b>	<b>377</b>
VI-1 Introduction . . . . .	377
VI-2 Annotated features . . . . .	378
VI-3 Sequences . . . . .	379
VI-4 Alignments . . . . .	384
<b>VII PLV Proteins</b>	<b>411</b>
VII-1 Introduction . . . . .	411
VII-2 Sequences . . . . .	412
VII-3 Alignments . . . . .	420



## I

# Preface

## I-1 Introduction

This compendium is an annual summary of the data contained in the HIV sequence database. We try to present a judicious selection of the data in such a way that it is of maximum utility to HIV researchers. Each of the alignments attempts to display the genetic variability within the different species, groups and subtypes of the virus.

This compendium contains sequences published before January 1, 2017. Hence, though it is published in 2017 and called the 2017 Compendium, its contents correspond to the 2016 curated alignments on our website.

The number of sequences in the HIV database is still increasing. In total, at the end of 2016, there were 748,589 sequences in the HIV Sequence Database, an increase of 10.8% since the previous year.

The number of near complete genomes (>7000 nucleotides) increased to 7416 by end of 2016. However, as in previous years, the compendium alignments contain only a fraction of these. A more complete version of all alignments is available on our website, <https://www.hiv.lanl.gov/content/sequence/NEWALIGN/align.html>

As always, we are open to complaints and suggestions for improvement. Inquiries and comments regarding the compendium should be addressed to [seq-info@lanl.gov](mailto:seq-info@lanl.gov).

## I-2 Acknowledgements

The HIV Sequence Database and Analysis Project is funded by the Vaccine and Prevention Research Program of the AIDS Division of the National Institute of Allergy and Infectious Diseases (Anjali Singh, Contracting Officer's Representative) through interagency agreement AAI 12007-0000-01000 "HIV/SIV Database and Analysis Unit" with the U.S. Department of Energy.

## I-3 Citing the compendium or database


The LANL HIV Sequence Database may be cited in the same manner as this compendium:

*HIV Sequence Compendium 2017*. Brian Foley, Thomas Leitner, Cristian Apetrei, Beatrice Hahn, Ilene Mizrahi, James Mullins, Andrew Rambaut, Steven Wolinsky, and Bette Korber editors. 2017. Publisher: Los Alamos National Laboratory, Theoretical Biology and Biophysics, Los Alamos, New Mexico. LA-UR-17-25240.

## I-4 About the PDF

The complete *HIV Sequence Compendium 2017: Preface* is available in Adobe Portable Document Format (PDF) from our website, <https://www.hiv.lanl.gov/>. The PDF version is hypertext enabled and features 'clickable' table-of-contents, indexes, references and links to external web sites. This volume is typeset using L<sup>A</sup>T<sub>E</sub>X.

I-5 Genome maps



**Landmarks of the HIV-1, HIV-2, and SIV genomes.** Open reading frames are shown as rectangles. The gene start, indicated by the small number in the upper left corner of each rectangle normally records the position of the **a** in the **atg** start codon for that gene, while the number in the lower right records the last position of the stop codon. For *pol*, the start is taken to be the first **t** in the sequence **tttttag**, which forms part of the stem loop that potentiates ribosomal slippage on the RNA and a resulting  $-1$  frameshift and the translation of the Gag-Pol polyprotein. The *tat* and *rev* spliced exons are shown as shaded rectangles. In HXB2, \*5772 marks the position of a frameshift in the *vpr* gene caused by an “extra” **t** relative to most other subtype B viruses; !6062 indicates a defective **acg** start codon in *vpu*; †8424 and †9168 mark premature stop codons in *tat* and *nef*. See Korber *et al.*, Numbering Positions in HIV Relative to HXB2CG, in *Human Retroviruses and AIDS*, 1998, p. 102. Available from <https://www.hiv.lanl.gov/content/sequence/HIV/REVIEWS/HXB2.html>

## I-6 HIV/SIV proteins

Name	Size	Function	Localization
<b>Gag</b>			
MA	p17	membrane anchoring; env interaction; nuclear transport of viral core (myristylated protein)	virion
CA	p24	core capsid	virion
NC	p7	nucleocapsid, binds RNA	virion
	p6	binds Vpr	virion
<b>Pol</b>			
Protease (PR)	p15	Gag/Pol cleavage and maturation	virion
Reverse Transcriptase (RT)	p66, p51	reverse transcription, RNase H activity	virion
RNase H	p15		virion
Integrase (IN)	p31	DNA provirus integration	virion
Env	gp120/gp41	external viral glycoproteins bind to CD4 and secondary receptors	plasma membrane, virion envelope
Tat	p16/p14	viral transcriptional transactivator	primarily in nucleolus/nucleus
Rev	p19	RNA transport, stability and utilization factor (phosphoprotein)	primarily in nucleolus/nucleus shuttling between nucleolus and cytoplasm
Vif	p23	promotes virion maturation and infectivity	cytoplasm (cytosol, membranes), virion
Vpr	p10-15	promotes nuclear localization of preintegration complex, inhibits cell division, arrests infected cells at G2/M	virion nucleus (nuclear membrane?)
Vpu	p16	promotes extracellular release of viral particles; degrades CD4 in the ER; (phosphoprotein only in HIV-1 and SIVcpz)	integral membrane protein
Nef	p27-p25	CD4 and class I downregulation (myristylated protein)	plasma membrane, cytoplasm, (virion?)
Vpx	p12-16	Vpr homolog present in HIV-2 and some SIVs, absent in HIV-1	virion (nucleus?)
Tev	p28	tripartite tat-env-rev protein (also named Tnv)	primarily in nucleolus/nucleus

## I-7 Landmarks of the genome

### HIV genomic structural elements

**LTR** Long terminal repeat, the DNA sequence flanking the genome of integrated proviruses. It contains important regulatory regions, especially those for transcription initiation and polyadenylation.

**TAR** Target sequence for viral transactivation, the binding site for Tat protein and for cellular proteins; consists of approximately the first 45 nucleotides of the viral mRNAs in HIV-1 (or the first 100 nucleotides in HIV-2 and SIV.) TAR RNA forms a hairpin stem-loop structure with a side bulge; the bulge is necessary for Tat binding and function.

**RRE** Rev responsive element, an RNA element encoded within the env region of HIV-1. It consists of approximately 200 nucleotides (positions 7327 to 7530 from the start of transcription in HIV-1, spanning the border of gp120 and gp41). The RRE is necessary for Rev function; it contains a high affinity site for Rev; in all, approximately seven binding sites for Rev exist within the RRE RNA. Other lentiviruses (HIV-2, SIV, visna, CAEV) have similar RRE elements in similar locations within env, while HTLVs have an analogous RNA element (RXRE) serving the same purpose within their LTR; RRE is the binding site for Rev protein, while RXRE is the binding site for Rex protein. RRE (and RXRE) form complex secondary structures, necessary for specific protein binding.

**PE** Psi elements, a set of 4 stem-loop structures preceding and overlapping the Gag start codon which are the sites recognized by the cysteine histidine box, a conserved motif with the canonical sequence CysX2CysX4HisX4Cys, present in the Gag p7 NC protein. The Psi Elements are present in unspliced genomic transcripts but absent from spliced viral mRNAs.

**SLIP** A TTTTTT slippery site, followed by a stem-loop structure, is responsible for regulating the -1 ribosomal frameshift out of the Gag reading frame into the Pol reading frame.

**CRS** Cis-acting repressive sequences postulated to inhibit structural protein expression in the absence of Rev. One such site was mapped within the pol region of HIV-1. The exact function has not been defined; splice sites have been postulated to act as CRS sequences.

**INS** Inhibitory/Instability RNA sequences found within the structural genes of HIV-1 and of other complex retroviruses. Multiple INS elements exist within the genome and can act independently; one of the best characterized elements spans nucleotides 414 to 631 in the gag region of HIV-1. The INS elements have been defined by functional assays as elements that inhibit expression posttranscriptionally. Mutation of the RNA elements was shown to lead to INS inactivation and up regulation of gene expression.

### Genes and gene products

**GAG** The genomic region encoding the capsid proteins (group specific antigens). The precursor is the p55 myristylated pro-

tein, which is processed to p17 (MA<sub>TR</sub>ix), p24 (CA<sub>PS</sub>id), p7 (Nucleo<sub>CA</sub>psid), and p6 proteins, by the viral protease. Gag associates with the plasma membrane where the virus assembly takes place. The 55 kDa Gag precursor is called assemblin to indicate its role in viral assembly.

**POL** The genomic region encoding the viral enzymes protease, reverse transcriptase, RNase, and integrase. These enzymes are produced as a Gag-Pol precursor polyprotein, which is processed by the viral protease; the Gag-Pol precursor is produced by ribosome frameshifting near the 3' end of gag.

**ENV** Viral glycoproteins produced as a precursor (gp160) which is processed to give a noncovalent complex of the external glycoprotein gp120 and the transmembrane glycoprotein gp41. The mature gp120-gp41 proteins are bound by non-covalent interactions and are associated as a trimer on the cell surface. A substantial amount of gp120 can be found released in the medium. gp120 contains the binding site for the CD4 receptor, and the seven transmembrane domain chemokine receptors that serve as co-receptors for HIV-1.

**TAT** Transactivator of HIV gene expression. One of two essential viral regulatory factors (Tat and Rev) for HIV gene expression. Two forms are known, Tat-1 exon (minor form) of 72 amino acids and Tat-2 exon (major form) of 86 amino acids. Low levels of both proteins are found in persistently infected cells. Tat has been localized primarily in the nucleolus/nucleus by immunofluorescence. It acts by binding to the TAR RNA element and activating transcription initiation and elongation from the LTR promoter, preventing the 5'LTR AATAAA polyadenylation signal from causing premature termination of transcription and polyadenylation. It is the first eukaryotic transcription factor known to interact with RNA rather than DNA and may have similarities with prokaryotic anti-termination factors. Extracellular Tat can be found and can be taken up by cells in culture.

**REV** The second necessary regulatory factor for HIV expression. A 19 kDa phosphoprotein, localized primarily in the nucleolus/nucleus, Rev acts by binding to RRE and promoting the nuclear export, stabilization and utilization of the unspliced viral mRNAs containing RRE. Rev is considered the most functionally conserved regulatory protein of lentiviruses. Rev cycles rapidly between the nucleus and the cytoplasm.

**VIF** Viral infectivity factor, a basic protein of typically 23 kDa. Promotes the infectivity but not the production of viral particles. In the absence of Vif the produced viral particles are defective, while the cell-to-cell transmission of virus is not affected significantly. Found in almost all lentiviruses, Vif is a cytoplasmic protein, existing in both a soluble cytosolic form and a membrane-associated form. The latter form of Vif is a peripheral membrane protein that is tightly associated with the cytoplasmic side of cellular membranes. In 2003, it was discovered that Vif prevents the action of the cellular APOBEC-3G protein which deaminates DNA:RNA heteroduplexes in the cytoplasm.

**VPR** Vpr (viral protein R) is a 96-amino acid (14 kDa) protein, which is incorporated into the virion. It interacts with the p6



Gag part of the Pr55 Gag precursor. Vpr detected in the cell is localized to the nucleus. Proposed functions for Vpr include the targeting the nuclear import of preintegration complexes, cell growth arrest, transactivation of cellular genes, and induction of cellular differentiation. In HIV-2, SIV-SMM, SIV-RCM, SIV-MND-2 and SIV-DRL the Vpx gene is apparently the result of a Vpr gene duplication event, possibly by recombination.

**VPU** Vpu (viral protein U) is unique to HIV-1, SIVcpz (the closest SIV relative of HIV-1), SIV-GSN, SIV-MUS, SIV-MON and SIV-DEN. There is no similar gene in HIV-2, SIV-SMM or other SIVs. Vpu is a 16 kDa (81-amino acid) type I integral membrane protein with at least two different biological functions: (a) degradation of CD4 in the endoplasmic reticulum, and (b) enhancement of virion release from the plasma membrane of HIV-1-infected cells. Env and Vpu are expressed from a bicistronic mRNA. Vpu probably possesses an N-terminal hydrophobic membrane anchor and a hydrophilic moiety. It is phosphorylated by casein kinase II at positions Ser52 and Ser56. Vpu is involved in Env maturation and is not found in the virion. Vpu has been found to increase susceptibility of HIV-1 infected cells to Fas killing.

**NEF** A multifunctional 27-kDa myristylated protein produced by an ORF located at the 3' end of the primate lentiviruses. Other forms of Nef are known, including nonmyristylated variants. Nef is predominantly cytoplasmic and associated with the plasma membrane via the myristyl residue linked to the conserved second amino acid (Gly). Nef has also been identified in the nucleus and found associated with the cytoskeleton in some experiments. One of the first HIV proteins to be produced in infected cells, it is the most immunogenic of the accessory proteins. The nef genes of HIV and SIV are dispensable *in vitro*, but are essential for efficient viral spread and disease progression *in vivo*. Nef is necessary for the maintenance of high virus loads and for the development of AIDS in macaques, and viruses with defective Nef have been detected in some HIV-1 infected long term survivors. Nef downregulates CD4, the primary viral receptor, and MHC class I molecules, and these functions map to different parts of the protein. Nef interacts with components of host cell signal transduction and clathrin-dependent protein sorting pathways. It increases viral infectivity. Nef contains PXXP motifs that bind to SH3 domains of a subset of Src kinases and are required for the enhanced growth of HIV but not for the downregulation of CD4.

**VPX** A virion protein of 12 kDa found in HIV-2, SIV-SMM, SIV-RCM, SIV-MND-2 and SIV-DRL and not in HIV-1 or other SIVs. This accessory gene is a homolog of HIV-1 vpr, and viruses with Vpx carry both vpr and vpx. Vpx function in relation to Vpr is not fully elucidated; both are incorporated into virions at levels comparable to Gag proteins through interactions with Gag p6. Vpx is necessary for efficient replication of SIV-SMM in PBMCs. Progression to AIDS and death in SIV-infected animals can occur in the absence of Vpr or Vpx. Double mutant virus lacking both vpr and vpx was at-

tenuated, whereas the single mutants were not, suggesting a redundancy in the function of Vpr and Vpx related to virus pathogenicity.

**ASP** In 1988, a study by Roger Miller noted the existence of an antisense open reading frame in HIV-1 that encodes an Antisense Protein (ASP) of 190 amino acids. The open reading frame occurs in the env region of the genome, which undergoes strong selection for divergence. Since many mutations that might occur in Env introduce stop codons in alternate reading frames, the high retention of the ASP open reading frame suggests a role for the product. The open reading frame is absent in most SIVcpz and O group sequences, but it is present in the majority of HIV-1 M group sequences.

**Structural proteins/viral enzymes** The products of *gag*, *pol*, and *env* genes, which are essential components of the retroviral particle.

**Regulatory proteins** Tat and Rev proteins of HIV/SIV and Tax and Rex proteins of HTLVs. They modulate transcriptional and posttranscriptional steps of virus gene expression and are essential for virus propagation.

**Accessory or auxiliary proteins** Additional virion and non-virion-associated proteins produced by HIV/SIV retroviruses: Vif, Vpr, Vpu, Vpx, Nef. Although the accessory proteins are in general not necessary for viral propagation in tissue culture, they have been conserved in the different isolates; this conservation and experimental observations suggest that their role *in vivo* is very important. Their functional importance continues to be elucidated.

**Complex retroviruses** Retroviruses regulating their expression via viral factors and expressing additional proteins (regulatory and accessory) essential for their life cycle.

**I-8 Amino acid codes**

---

A	Alanine
B	Aspartic Acid or Asparagine
C	Cysteine
D	Aspartic Acid
E	Glutamic Acid
F	Phenylalanine
G	Glycine
H	Histidine
I	Isoleucine
K	Lysine
L	Leucine
M	Methionine
N	Asparagine
P	Proline
Q	Glutamine
R	Arginine
S	Serine
T	Threonine
V	Valine
W	Tryptophan
X	unknown or "other" amino acid
Y	Tyrosine
Z	Glutamic Acid or Glutamine
.	gap
-	identity
*	stop codon
#	incomplete codon

---

**I-9 Nucleic acid codes**

---

A	Adenine
C	Cytosine
G	Guanine
T	Thymine
U	Uracil
M	A or C
R	A or G
W	A or T
S	C or G
Y	C or T
K	G or T
V	A or C or G
H	A or C or T
D	A or G or T
B	C or G or T
N	unknown
.	gap
-	identity

---